

B6

(4)

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 October 2003 (02.10.2003)

PCT

(10) International Publication Number
WO 03/080573 A1

(51) International Patent Classification⁷: **C07D 209/48**,
405/06, 409/06, 413/06, 471/04, A61K 31/4035, 31/437 //
(C07D 471/04, 221/00, 209/00)

(21) International Application Number: PCT/EP03/02931

(22) International Filing Date: 20 March 2003 (20.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
02007222.9 27 March 2002 (27.03.2002) EP

(71) Applicant: **F. HOFFMANN-LA ROCHE AG** [CH/CH];
Grenzacherstrasse 124, CH-4070 Basle (CH).

(72) Inventors: **CESURA, Andrea**; 26, rue du Grand Pre,
CH-1299 Crans-pres-Celigny (CH). **RODRIGUEZ**
SARMIENTO, Rosa, Maria; Missionsstrasse 33,
CH-4055 Basel (CH). **THOMAS, Andrew, William**;
Zwinglistrasse 4, CH-4127 Birsfelden (CH). **WYLER,**
Rene; Brandschenkestrasse 168, CH-8002 Zuerich (CH).

(74) Agent: **WAECHTER, Dieter**; Grenzacherstrasse 124,
CH-4070 Basle (CH).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU,
ZA, ZM, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **PHTHALIMIDO DERIVATIVES AS INHIBITORS OF MONOAMINE OXIDASE B**




(57) **Abstract:** This invention relates to phthalimido derivatives of the general formula (I), wherein X is -N= or -CH, and R¹ to R⁴ and m are as defined in the specification, as well as their pharmaceutically acceptable salts. The invention further relates to medicaments containing these compounds, a process for their preparation as well as their use for preparation of medicaments for the treatment or prevention of diseases in which MAO-B inhibitors might be beneficial.

WO 03/080573 A1

PHTHALIMIDO DERIVATIVES AS INHIBITORS OF MONOAMINE OXIDASE B

This invention relates to phthalimido derivatives of the general formula



wherein

X is $-N=$ or $-CH=$;

- 5 R^1 is $-CO-NR^5R^6$; $-CHR^7-(CH_2)_n-CO-NR^5R^6$;
 $-(CH_2)_n-NR^5R^6$;
 $-(CH_2)_n-COOR^8$;
 $-(CH_2)_n-CN$;
 $-CHR^7-(CH_2)_n-CF_3$;
10 $-(CH_2)_n-NH-COR^9$;
 $-(CH_2)_n-NH-COOR^8$;
 $-(CH_2)_n$ -piperidinyl, $-(CH_2)_n$ -morpholinyl, $-(CH_2)_n$ -tetrahydrofuranyl;
 $-(CH_2)_n$ -thiophenyl or $-(CH_2)_n$ -isoxazolyl, wherein the heterocyclic ring may
be substituted by C_1-C_6 -alkyl;
15 $-(CH_2)_n$ -phenyl, wherein the phenyl ring may be substituted by halogen or
halogen- (C_1-C_6) -alkyl;
 $-(CH_2)_p-OR^8$;
 $-(CH_2)_p-SR^8$;
 $-(CH_2)_p-SO-R^9$; or
20 $-(CH_2)_n-CS-NR^5R^6$;
- R^2 is hydrogen; C_1-C_6 -alkyl;
 $-(CH_2)_p-OR^{10}$;
 $-(CH_2)_p-SR^{10}$; or benzyl;
- R^3 is hydrogen or C_1-C_6 -alkyl;

- R^4 is halogen, halogen-(C_1-C_6)-alkyl, cyano, C_1-C_6 -alkoxy or halogen-(C_1-C_6)-alkoxy;
- R^5 and R^6 are independently from each other hydrogen or C_1-C_6 -alkyl;
- R^7 is hydrogen, hydroxy or C_1-C_6 -alkoxy;
- 5 R^8 is hydrogen or C_1-C_6 -alkyl;
- R^9 is C_1-C_6 -alkyl;
- R^{10} is hydrogen or C_1-C_6 -alkyl;
- m is 1, 2 or 3;
- n is 0, 1 or 2; and
- 10 p is 1 or 2;

as well as their pharmaceutically acceptable salts.

It has now been found that the compounds of general formula I are selective monoamine oxidase B inhibitors.

- Monoamine oxidase (MAO, EC 1.4.3.4) is a flavin-containing enzyme responsible
- 15 for the oxidative deamination of endogenous monoamine neurotransmitters such as dopamine, serotonin, adrenaline, or noradrenaline, and trace amines, e.g. phenylethylamine, as well as a number of amine xenobiotics. The enzyme exists in two forms, MAO-A and MAO-B, encoded by different genes (A. W. Bach et al., *Proc. Natl. Acad. Sci. USA* 1988, 85, 4934-4938) and differing in tissue distribution, structure and substrate
- 20 specificity. MAO-A has higher affinity for serotonin, octopamine, adrenaline, and noradrenaline; whereas the natural substrates for MAO-B are phenylethylamine and tyramine. Dopamine is thought to be oxidised by both isoforms. MAO-B is widely distributed in several organs including brain (A.M. Cesura and A. Pletscher, *Prog. Drug Research* 1992, 38, 171-297). Brain MAO-B activity appears to increase with age. This
- 25 increase has been attributed to the gliosis associated with aging (C.J. Fowler et al., *J. Neural. Transm.* 1980, 49, 1-20). Additionally, MAO-B activity is significantly higher in the brains of patients with Alzheimer's disease (P. Dostert et al., *Biochem. Pharmacol.* 1989, 38, 555-561) and it has been found to be highly expressed in astrocytes around senile plaques (Saura et al., *Neuroscience* 1994, 70, 755-774). In this context, since
- 30 oxidative deamination of primary monoamines by MAO produces NH_3 , aldehydes and H_2O_2 , agents with established or potential toxicity, it is suggested that there is a rationale for the use of selective MAO-B inhibitors for the treatment of dementia and Parkinson's disease. Inhibition of MAO-B causes a reduction in the enzymatic inactivation of dopamine and thus prolongation of the availability of the neurotransmitter in
- 35 dopaminergic neurons. The degeneration processes associated with age and Alzheimer's

and Parkinson's diseases may also be attributed to oxidative stress due to increased MAO activity and consequent increased formation of H_2O_2 by MAO-B. Therefore, MAO-B inhibitors may act by both reducing the formation of oxygen radicals and elevating the levels of monoamines in the brain.

5 Given the implication of MAO-B in the neurological disorders mentioned above, there is considerable interest to obtain potent and selective inhibitors that would permit control over this enzymatic activity. The pharmacology of some known MAO-B inhibitors is for example discussed by D. Bentué-Ferrer et al. in *CNS Drugs* 1996, 6, 217-236. Whereas a major limitation of irreversible and non-selective MAO inhibitor activity
10 is the need to observe dietary precautions due to the risk of inducing a hypertensive crisis when dietary tyramine is ingested, as well as the potential for interactions with other medications (D. M. Gardner et al., *J. Clin. Psychiatry* 1996, 57, 99-104), these adverse events are of less concern with reversible and selective MAO inhibitors, in particular of MAO-B. Thus, there is a need for MAO-B inhibitors with a high selectivity and without
15 the adverse side-effects typical of irreversible MAO inhibitors with low selectivity for the enzyme.

 The object of the present invention therefore is to provide compounds which must have the advantageous properties mentioned above. It has been found that the compounds of formula I of the present invention and their pharmaceutically acceptable
20 salts show the potential to be highly selective MAO-B inhibitors. Subjects of the present invention are further medicaments based on a compound of formula I in accordance with the invention, a process for the manufacture of compounds of formula I and their pharmaceutically acceptable salts, as well as the use of the compounds of formula I in the control or prevention of diseases mediated by monoamine oxidase B inhibitors, and,
25 respectively, the use for the production of corresponding medicaments.

 The following definitions of general terms used in the present patent application apply irrespective of whether the terms in question appear alone or in combination. It must be noted that, as used in the specification and the appended claims, the singular forms "a", "an," and "the" include plural forms unless the context clearly dictates
30 otherwise.

 The term "C₁-C₆-alkyl" ("lower alkyl") used in the present application denotes straight-chain or branched saturated hydrocarbon residues with 1 to 6 carbon atoms, preferably with 1 to 4 carbon atoms, such as methyl, ethyl, n-propyl, i-propyl, n-butyl, sec-butyl, t-butyl, and the like.

35 The term "halogen" denotes fluorine, chlorine, bromine and iodine.

"Halogen-(C₁-C₆)-alkyl" or "halogen-(C₁-C₆)-alkoxy" means the lower alkyl residue or lower alkoxy residue, respectively, as defined herein substituted in any position with one or more halogen atoms as defined herein. Examples of halogenalkyl residues include, but are not limited to, 1,2-difluoropropyl, 1,2-dichloropropyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2,2,2-trichloroethyl, and 1,1,1-trifluoropropyl, and the like. "Halogenalkoxy" includes trifluoromethoxy.

"C₁-C₆-Alkoxy" means the residue -O-R, wherein R is a lower alkyl residue as defined herein. Examples of alkoxy radicals include, but are not limited to, methoxy, ethoxy, isopropoxy, and the like.

10 "Pharmaceutically acceptable salts" of a compound means salts that are pharmaceutically acceptable, which are generally safe, non-toxic, and neither biologically nor otherwise undesirable, and that possess the desired pharmacological activity of the parent compound. These salts are derived from an inorganic or organic acid or base.

Such salts include:

- 15 (1) acid addition salts formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, benzenesulfonic acid, benzoic, camphorsulfonic acid, citric acid, ethanesulfonic acid, fumaric acid, glucoheptonic acid, gluconic acid, glutamic acid, glycolic acid, hydroxynaphthoic acid, 2-
20 hydroxyethanesulfonic acid, lactic acid, maleic acid, malic acid, mandelic acid, methanesulfonic acid, muconic acid, 2-naphthalenesulfonic acid, propionic acid, salicylic acid, succinic acid, dibenzoyl-L-tartaric acid, tartaric acid, p-toluene-sulfonic acid, trimethylacetic acid, 2,2,2-trifluoroacetic acid, and the like; or
- (2) salts formed when an acidic proton present in the parent compound either is replaced
25 by a metal ion, e.g., an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic or inorganic base. Acceptable organic bases include diethanolamine, ethanolamine, N-methylglucamine, triethanolamine, tromethamine, and the like. Acceptable inorganic bases include aluminum hydroxide, calcium hydroxide, potassium hydroxide, sodium carbonate and sodium hydroxide.

30 It should be understood that all references to pharmaceutically acceptable salts include solvent addition forms (solvates) or crystal forms (polymorphs) of the same acid addition salt.

Among compounds of the present invention certain compounds of formula I, or pharmaceutically acceptable salts thereof, are preferred.

Preferred compounds of formula I are those, in which X is $-\text{CH}=\text{}$.

Especially preferred are compounds of formula I, in which R^1 is $-\text{CO}-\text{NR}^5\text{R}^6$ or $-\text{CHR}^7-(\text{CH}_2)_n-\text{CO}-\text{NR}^5\text{R}^6$, and wherein R^5 and R^6 are independently from each other hydrogen or C_1 - C_6 -alkyl, R^7 is hydrogen, hydroxy or C_1 - C_6 -alkoxy and n is 0, 1 or 2.

5 Examples of such compounds are the following:

- 2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetamide,
 (S)-2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-propionamide,
 (S)-2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-3-hydroxy-
 propionamide,
 10 (R)-2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-propionamide,
 2-[5-(3-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-propionamide,
 (2-[5-(3-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetamide, and
 2-[5-(3-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-3-hydroxy-
 propionamide.

15 Also preferred are compounds of formula I, in which R^1 is $-(\text{CH}_2)_n-\text{NR}^5\text{R}^6$, $-(\text{CH}_2)_n-\text{NH}-\text{COR}^9$ or $-(\text{CH}_2)_n$ -piperidinyl, and wherein R^5 and R^6 are independently from each other hydrogen or C_1 - C_6 -alkyl, R^9 is C_1 - C_6 -alkyl and n is 0, 1, or 2.

The following compounds are examples thereof:

- N-[2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-ethyl]-acetamide,
 20 2-(2-amino-ethyl)-5-(4-fluoro-benzyloxy)-isoindole-1,3-dione, and
 5-(4-fluoro-benzyloxy)-2-piperidin-4-yl-isoindole-1,3-dione.

Further preferred compounds of formula I are those, in which R^1 is $-(\text{CH}_2)_p-\text{OR}^8$ or $-\text{CHR}^7-(\text{CH}_2)_n-\text{CF}_3$, and wherein R^7 is hydrogen, hydroxy or C_1 - C_6 -alkoxy, R^8 is hydrogen or C_1 - C_6 -alkyl and p is 1 or 2.

25 Examples of such compounds are the following:

- 5-(4-fluoro-benzyloxy)-2-(2-hydroxy-ethyl)-isoindole-1,3-dione,
 5-(4-fluoro-benzyloxy)-2-(2-methoxy-ethyl)-isoindole-1,3-dione,
 5-(3-fluoro-benzyloxy)-2-(2-methoxy-ethyl)-isoindole-1,3-dione,
 (S)-5-(4-fluoro-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione,
 30 (S)-5-(3-fluoro-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione,
 (S)-5-(2-fluoro-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione,
 (S)-2-(2-methoxy-1-methyl-ethyl)-5-(4-trifluoromethyl-benzyloxy)-isoindole-1,3-dione,
 (S)-5-(4-bromo-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione,
 (S)-5-(3,4-difluoro-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione,
 35 5-(3-fluoro-benzyloxy)-2-(2-hydroxy-ethyl)-isoindole-1,3-dione,
 5-(4-fluoro-benzyloxy)-2-(3,3,3-trifluoro-2-hydroxy-propyl)-isoindole-1,3-dione, and

5-(3,5-bis-trifluoromethyl-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione.

Preferred compounds of formula I are further those, in which R^1 is $-(CH_2)_p-SR^8$; $-(CH_2)_p-SO-R^9$; or $-(CH_2)_n-CS-NR^5R^6$, and wherein R^5 and R^6 are independently from each other hydrogen or C_1-C_6 -alkyl, R^8 is hydrogen or C_1-C_6 -alkyl, R^9 is C_1-C_6 -alkyl and n is 0, 1, or 2.

Examples of such compounds are the following:

2-(2-ethylsulfanyl-ethyl)-5-(4-fluoro-benzyloxy)-isoindole-1,3-dione,
(S)-2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-thiopropionamide,
10 and
2-(2-ethylsulfanyl-ethyl)-5-(3-fluoro-benzyloxy)-isoindole-1,3-dione.


Also preferred are compounds of formula I, in which R^1 is $-(CH_2)_n-CN$ and n is 0, 1 or 2.

The following compounds are examples thereof:

15 5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetonitrile, and
[5-(3-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetonitrile.


Preferred are also compounds of formula I, in which R^4 signifies halogen. Especially preferred are compounds of formula I, in which R^4 is fluoro and m is 1.

The compounds of general formula I and their pharmaceutically acceptable salts
20 can be manufactured by reacting a compound of formula



II


with a compound of formula



III

to obtain a compound of formula


- 7 -



I


and, if desired, converting a compound of formula I into a pharmaceutically acceptable salt.

Alternatively, the compounds of general formula I and their pharmaceutically acceptable salts can be manufactured by reacting a compound of formula




IV

with a compound of formula



V

to obtain a compound of formula




I

and, if desired, converting a compound of formula I into a pharmaceutically acceptable salt.

10


In accordance with the present invention, compounds of general formula I can be prepared following scheme 1: A compound of formula VI is heated in the presence of ammonium carbonate. The obtained compound VII is then treated with benzylic bromides in the presence of potassium carbonate to afford compounds of type IV which are then dissolved in THF and treated with sodium hydride and an electrophile of formula V to give compounds of formula I.

Scheme 1



Alternatively, in accordance with the present invention, compounds of general formula I can be prepared following scheme 2: A compound of formula VI is heated in the presence of benzylic bromides and potassium carbonate to afford VIII which are then saponified with bases such as LiOH to give IX. Treatment of IX with an activating agent such as carbonyldiimidazole in an appropriate solvent such as DMF or DMA or NMP, followed by heating (conventional or microwave) in the presence of amines III affords compounds of formula I.


Scheme 2



Alternatively, in accordance with the present invention, compounds of general formula I can be prepared following scheme 3: A compound of formula VI is heated in the presence of an amine III with an activating agent such as carbonyldiimidazole in an appropriate solvent such as dimethylformamide (DMF) or dimethylacetamide (DMA) or N-methylpyrrolidone (NMP), followed by heating (conventional or microwave) to afford compounds of formula X which are in turn reacted with benzyl bromides in the presence of potassium carbonate to give compounds of formula I.


10

Scheme 3



Alternatively, in accordance with the present invention, compounds of general formula XVI can be prepared following scheme 4: A compound of formula XI is oxidised to XII and chlorinated to give XIII via successive treatment with m-chloroperoxybenzoic acid (m-CPBA) followed by phosphorus oxychloride. Subsequent reaction with sodium salts of benzyl alcohols affords XIV which after saponification to compounds of type XV are treated with an activating agent such as carbonyldiimidazole in an appropriate solvent such as DMF or DMA or NMP, followed by heating (conventional or microwave) in the presence of amines III affords compounds of formula XVI.

Scheme 4



- Pharmaceutically acceptable salts of compounds of formula I can be manufactured readily according to methods known per se and taking into consideration the nature of the compound to be converted into a salt. Inorganic or organic acids such as, for example, hydrochloric acid, hydrobromic acid, sulphuric acid, nitric acid, phosphoric acid or citric acid, formic acid, fumaric acid, maleic acid, acetic acid, succinic acid, tartaric acid, methanesulphonic acid, p-toluenesulphonic acid and the like are suitable for the formation of pharmaceutically acceptable salts of basic compounds of formula I.
- Compounds which contain the alkali metals or alkaline earth metals, for example sodium, potassium, calcium, magnesium or the like, basic amines or basic amino acids are suitable for the formation of pharmaceutically acceptable salts of acidic compounds.

- The compounds of formula I and their pharmaceutically acceptable salts are, as already mentioned above, monoamine oxidase B inhibitors and can be used for the treatment or prevention of diseases in which MAO-B inhibitors might be beneficial. These include acute and chronic neurological disorders, cognitive disorders and memory deficits. Treatable neurological disorders are for instance traumatic or chronic degenerative processes of the nervous system, such as Alzheimer's disease, other types of

dementia, minimal cognitive impairment or Parkinson's disease. Other indications include psychiatric diseases such as depression; anxiety, panic attack, social phobia, schizophrenia, eating and metabolic disorders such as obesity as well as the prevention and treatment of withdrawal syndromes induced by abuse of alcohol, nicotine and other
5 addictive drugs. Other treatable indications may be reward deficiency syndrome (G.M. Sullivan, International patent application No. WO 01/34172 A2), peripheral neuropathy caused by cancer chemotherapy (G. Bobotas, International Patent Application No. WO 97/33572 A1), or the treatment of multiple sclerosis (R.Y. Harris, International patent application No. WO 96/40095 A1) and other neuroinflammatory diseases.

10 The compounds of formula I and their pharmaceutically acceptable salts are especially useful for the treatment and prevention of Alzheimer's disease and senile dementia.

The pharmacological activity of the compounds was tested using the following method:

15 The cDNA's encoding human MAO-A and MAO-B were transiently transfected into EBNA cells using the procedure described by E.-J. Schlaeger and K. Christensen (Transient Gene Expression in Mammalian Cells Grown in Serum-free Suspension Culture; Cytotechnology, 15: 1-13, 1998). After transfection, cells were homogenised by means of a Polytron homogenizer in 20 mM Tris HCl buffer, pH 8.0, containing 0.5 mM
20 EGTA and 0.5 mM phenylmethanesulfonyl fluoride. Cell membranes were obtained by centrifugation at 45,000 x g and, after two rinsing step with 20 mM Tris HCl buffer, pH 8.0, containing 0.5 mM EGTA, membranes were eventually re-suspended in the above buffer and aliquots stored at -80 °C until use.

MAO-A and MAO-B enzymatic activity was assayed in 96-well-plates using a
25 spectrophotometric assay adapted from the method described by M. Zhou and N. Panchuk-Voloshina (A One-Step Fluorometric Method for the Continuous Measurement of Monoamine Oxidase Activity, Analytical Biochemistry, 253: 169-174, 1997). Briefly, membrane aliquots were incubated in 0.1 M potassium phosphate buffer, pH 7.4, for 30 min at 37 °C with or without various concentrations of the compounds.
30 After this period, the enzymatic reaction was started by the addition of the MAO substrate tyramine together with 1 U/ml horse-radish peroxidase (Roche Biochemicals) and 80 µM N-acetyl-3,7,-dihydroxyphenoxazine (Amplex Red, Molecular Probes). The samples were further incubated for 30 min at 37 °C in a final volume of 200 µl and absorbance was then determined at a wavelength of 570 nm using a SpectraMax plate
35 reader (Molecular Devices). Background (non-specific) absorbance was determined in the presence of 10 µM clorgyline for MAO-A or 10 µM L-deprenyl for MAO-B.

IC₅₀ values were determined from inhibition curves obtained using nine inhibitor concentrations in duplicate, by fitting data to a four parameter logistic equation using a computer program.

The compounds of the present invention are specific MAO-B inhibitors. The IC₅₀ values of compounds of formula I as measured in the assay described above are in the range of 10 µM or less, typically of 1 µM or less, ideally 0.03 µM or less, and more preferably 0.1 µM or less.

In the table below are described some specific IC₅₀ values of preferred compounds.

Compound	IC ₅₀ MAO-B (µM)	IC ₅₀ MAO-A (µM)
(S)-2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-propionamide (Example 2)	0.008	0.776
5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetonitrile (Example 7)	0.034	> 10
2-(2-Amino-ethyl)-5-(4-fluoro-benzyloxy)-isoindole-1,3-dione 1:1 hydrochloride (Example 9)	0.032	> 10
5-(4-Fluoro-benzyloxy)-2-(2-hydroxy-ethyl)-isoindole-1,3-dione (Example 12)	0.004	0.369
(S)-2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-3-hydroxy-propionamide (Example 14)	0.017	> 10
2-(2-Ethylsulfanyl-ethyl)-5-(4-fluoro-benzyloxy)-isoindole-1,3-dione (Example 19)	0.011	> 10
4-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-3-hydroxy-butyramide (Example 22)	0.097	> 10
(S)-5-(4-Bromo-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione (Example 28)	0.034	> 10
(S)-5-(3-Methoxy-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione (Example 29)	0.069	> 10
(S)-4-[2-(2-Methoxy-1-methyl-ethyl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-ylloxymethyl]-benzonitrile (Example 30)	0.129	6.200
(S)-5-(3,4-Difluoro-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione (Example 31)	0.032	3.496

Compound	IC ₅₀ MAO-B (μM)	IC ₅₀ MAO-A (μM)
(S)-2-(2-Methoxy-1-methyl-ethyl)-5-(4-trifluoromethoxy-benzyloxy)-isoindole-1,3-dione (Example 32)	0.047	> 10
5-(4-Fluoro-benzyloxy)-2-thiophen-2-ylmethyl-isoindole-1,3-dione (Example 34)	0.081	> 10
2-(2-Ethanesulfinyl-ethyl)-5-(4-fluoro-benzyloxy)-isoindole-1,3-dione (Example 38)	0.115	> 10
(5-(4-Fluoro-benzyloxy)-2-(3,3,3-trifluoro-2-methoxy-propyl)-isoindole-1,3-dione (Example 43)	0.045	> 10
(S)-2-[5-(3-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-propionamide (Example 44)	0.007	5.762
2-(2-Amino-ethyl)-5-(3-fluoro-benzyloxy)-isoindole-1,3-dione 1:1 hydrochloride (Example 50)	0.039	> 10
(S)-2-[5-(3-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-3-hydroxy-propionamide (Example 52)	0.020	> 10

The compounds of formula I and pharmaceutically acceptable salts thereof can be used as medicaments, e.g. in the form of pharmaceutical preparations. The pharmaceutical preparations can be administered orally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions. However, the administration can also be effected rectally, e.g. in the form of suppositories, or parenterally, e.g. in the form of injection solutions.

The compounds of formula I and pharmaceutically acceptable salts thereof can be processed with pharmaceutically inert, inorganic or organic carriers for the production of pharmaceutical preparations. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts and the like can be used, for example, as such carriers for tablets, coated tablets, dragées and hard gelatine capsules. Suitable carriers for soft gelatine capsules are, for example, vegetable oils, waxes, fats, semi-solid and liquid polyols and the like; depending on the nature of the active substance no carriers are, however, usually required in the case of soft gelatine capsules. Suitable carriers for the production of solutions and syrups are, for example, water, polyols, sucrose, invert sugar, glucose and the like. Adjuvants, such as alcohols, polyols, glycerol, vegetable oils and the like, can be used for aqueous injection solutions of water-soluble salts of compounds of formula I,

but as a rule are not necessary. Suitable carriers for suppositories are, for example, natural or hardened oils, waxes, fats, semi-liquid or liquid polyols and the like.

In addition, the pharmaceutical preparations can contain preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts
5 for varying the osmotic pressure, buffers, masking agents or antioxidants. They may also contain other therapeutically valuable substances.

As mentioned earlier, medicaments containing a compound of formula I or pharmaceutically acceptable salts thereof and a therapeutically inert excipient are also an object of the present invention, as is a process for the production of such medicaments
10 which comprises bringing one or more compounds of formula I or pharmaceutically acceptable salts thereof and, if desired, one or more other therapeutically valuable substances into a galenical dosage form together with one or more therapeutically inert carriers.

The dosage can vary within wide limits and will, of course, be fitted to the
15 individual requirements in each particular case. In general, the effective dosage for oral or parenteral administration is between 0.01-20 mg/kg/day, with a dosage of 0.1-10 mg/kg/day being preferred for all of the indications described. The daily dosage for an adult human being weighing 70 kg accordingly lies between 0.7-1400 mg per day, preferably between 7 and 700 mg per day.

20 The following examples are provided for illustration of the invention. They should not be considered as limiting the scope of the invention, but merely as being representative thereof.

Example 1

2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetamide

25 a) 4-(4-Fluoro-benzyloxy)-phthalic acid bis-(4-fluoro-benzyl) ester

A mixture of 4-hydroxyphthalic acid (3.5 g, 19 mmol), potassium carbonate (23.9 g, 173 mmol) and 4-fluorobenzylbromide (32.7 g, 173 mmol) in acetone (100 mL) and water (50 mL) was heated under reflux for 72 h. After cooling to room temperature, the mixture was evaporated and the product extracted with ethyl acetate. The organic
30 extracts were then washed with brine, dried over sodium sulfate. Filtration and evaporation gave a residue which was purified by chromatography (SiO₂, hexane-EtOAc 4:1) to afford the title compound (7.8 g, 80%) as a colourless oil. MS: m/e = 507.4 (M+H⁺).

b) 4-(4-Fluoro-benzyloxy)-phthalic acid (Method A)

- A mixture of 4-(4-fluoro-benzyloxy)-phthalic acid bis-(4-fluoro-benzyl) ester (7.8 g, 15.4 mmol) lithium hydroxide monohydrate (1.8 g, 46.2 mmol) in tetrahydrofuran (60 mL) and water (60 mL) was stirred at room temperature for 72 h. The mixture was then
5 evaporated followed by acidification to pH 2 at 0 °C with concentrated HCl. The organic layer was then diluted with ethyl acetate, separated, washed with water and dried over sodium sulfate. Filtration and half evaporation gave a white suspension which was filtered off and washed with ether to afford the title compound (2.5 g, 56%) as a white solid.
- 10 MS: $m/e = 289.0$ ($M-H^+$).

c) 4-(4-Fluoro-benzyloxy)-phthalic acid (Method B)

- A mixture of 4-hydroxyphthalic acid (5.0 g, 27 mmol), potassium hydroxide (5.4 g, 30 mmol) and 4-fluorobenzylbromide (5.7 g, 30 mmol) in water (13 mL) was heated under reflux for 5 h. After cooling to room temperature, the mixture was washed with
15 ether. The aqueous phase was then acidified to pH 2 at 0 °C with concentrated HCl and then extracted with ethyl acetate. The organic extracts were then washed with water and dried over sodium sulfate. Filtration and evaporation gave a residue which was purified by chromatography (SiO_2 , $CHCl_3$: Acetone: AcOH: 90:9:1) to afford the title compound (2.3 g, 29%) as a colourless oil. MS: $m/e = 289.0$ ($M-H^+$).

20 d) 2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetamide

- A mixture of 4-(4-fluoro-benzyloxy)-phthalic acid (185 mg, 0.64 mmol) and carbonyldiimidazole (109 mg, 0.67 mmol) in 1-methyl-2-pyrrolidinone (4 mL) was stirred at room temperature for 10 min and then heated at 50 °C for 15 min. To this mixture was added glycine hydrochloride (78 mg, 0.71 mmol) and pyridine (50 mg,
25 0.63 mmol) and the resulting mixture heated at 175 or 200 °C with microwave radiation (Smith reactor) for 10 min. After cooling to room temperature, the mixture was diluted with water and the resulting precipitate was filtered off and purified by chromatography (SiO_2 , CH_2Cl_2 : 2N NH_3 /MeOH 9:1) to afford the title compound (146 mg, 69%) as a white solid. MS: $m/e = 329.3$. ($M+H^+$).

- 30 Alternatively, 2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetamide was prepared from 5-hydroxy-isoindole-1,3-dione.

e) 5-Hydroxy-isoindole-1,3-dione

A mixture of 4-hydroxyphthalic acid (5.0 g, 27.0 mmol), acetic acid (25 mL), and ammonium carbonate (5.3 g, 55 mmol) was heated at 120 °C for 45 min followed by heating at 160 °C for 2 h. After cooling to room temperature, the reaction mixture was
5 half-evaporated and then the reaction mixture was basified to pH 10 with 1 N NaOH followed by acidification to pH 5 at 0 °C with concentrated HCl. The resulting precipitate was filtered off, washed with water, dried at 60 °C under high vacuum over night, to afford the title compound (3.2 g, 71%) as an off-white crystalline solid.
MS: $m/e = 162.1$ ($M-H^+$).

10 f) 5-(4-Fluoro-benzyloxy)-isoindole-1,3-dione

A mixture of 5-hydroxy-isoindole-1,3-dione (200 mg, 1.0 mmol), potassium carbonate (178 mg, 1.05 mmol), 4-fluorobenzylbromide (204 mg, 1.05 mmol) in ethanol (5 mL) was heated at 80 °C overnight. After cooling to room temperature, the reaction mixture was filtered and evaporated. The residue was purified by chromatography (SiO_2 ,
15 heptane- CH_2Cl_2 2:3 then CH_2Cl_2 - 2N NH_3 -MeOH 99:1: to 95:5) to afford the title compound (127 mg, 38%) as an off-white solid. MS: $m/e = 271.1$ (M^+).

g) 2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetamide

A mixture of 5-(4-fluoro-benzyloxy)-isoindole-1,3-dione (100 mg, 0.37 mmol), sodium hydride (55% in mineral oil, 18 mg, 0.42 mmol) and 2-bromoacetamide (61 mg,
20 0.44 mmol) in dry tetrahydrofuran (5 mL) at 0 °C was stirred at room temperature for 3 h and then heated at 50 °C for 1 h. After cooling to 0 °C, water (2 mL) was added and the product extracted with ethyl acetate. The organic extracts were then washed with brine and dried over sodium sulfate. Filtration and evaporation gave a residue which was purified by chromatography (SiO_2 , heptane- CH_2Cl_2 2:3 then CH_2Cl_2 - 2N NH_3 -MeOH
25 99:1: to 4:1) to afford the title compound (78 mg, 65%) as a white solid. MS: $m/e = 329.3$ ($M+H^+$).

Example 2

(S)-2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-propionamide

As described for example 1d, 4-(4-fluoro-benzyloxy)-phthalic acid (185 mg, 0.64
30 mmol) was converted to the title compound (106 mg, 49%) (using H-alanine- NH_2 HCl instead of glycineamide hydrochloride) which was obtained as a white solid.
MS: $m/e = 343.3$ ($M+H^+$).

- 17 -

Example 3

(S)-2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-3-methyl-
butyramide

As described for example 1d, 4-(4-fluoro-benzyloxy)-phthalic acid (185 mg, 0.64
5 mmol) was converted to the title compound (115 mg, 48%) (using H-valine-NH₂ HCl
instead of glycineamide hydrochloride) which was obtained as a white solid.

MS: m/e = 371.3. (M+H⁺).

Example 4

(S)-2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-4-methyl-
10 pentanoic acid amide

As described for example 1d, 4-(4-fluoro-benzyloxy)-phthalic acid (185 mg, 0.64
mmol) was converted to the title compound (126 mg, 51%) (using H-leucine-NH₂ HCl
instead of glycineamide hydrochloride) which was obtained as a white solid.

MS: m/e = 385.3 (M+H⁺).

15

Example 5

(R,S)-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-4-methylsulfanyl-
butyramide

As described for example 1d, 4-(4-fluoro-benzyloxy)-phthalic acid (165 mg, 0.57
mmol) was converted to the title compound (42 mg, 18%) (using DL-methionamide-HCl
20 instead of glycineamide hydrochloride) which was obtained as an off-white foam.

MS: m/e = 403.4 (M+H⁺).

Example 6

[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetic acid ethyl ester

As described for example 1d, 4-(4-fluoro-benzyloxy)-phthalic acid (185 mg, 0.64
25 mmol) was converted to the title compound (110 mg, 48%) (using glycine ether ester-
HCl instead of glycineamide hydrochloride) which was obtained as a white solid.

MS: m/e = 358.2 (M+H⁺).

Example 7

5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetonitrile

As described for example 1d, 4-(4-fluoro-benzyloxy)-phthalic acid (300 mg, 1.0 mmol) was converted to the title compound (215 mg, 67%) (using aminoacetonitrile instead of glycineamide hydrochloride) which was obtained as a white solid.
MS: $m/e = 328.2$ ($M+NH_4^+$).

Example 8

N-{2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-ethyl}-acetamide

A mixture of 4-(4-fluoro-benzyloxy)-phthalic acid (185 mg, 0.64 mmol) and carbonyldiimidazole (109 mg, 0.67 mmol) in N,N-dimethylacetamide (3 mL) was stirred at room temperature for 10 min and then heated at 50 °C for 15 min. To this mixture was added N-acetylenediamine (78 mg, 0.76 mmol) and the resulting mixture heated under reflux for 20 min. After cooling to room temperature, the mixture was evaporated and the residue was purified by chromatography (SiO_2 , CH_2Cl_2 : 2N NH_3 /MeOH 98:2 to 9:1) to afford the title compound (139 mg, 62%) as an off-white solid.
MS: $m/e = 357.3$. ($M+H^+$).

Example 9

2-(2-Amino-ethyl)-5-(4-fluoro-benzyloxy)-isoindole-1,3-dione 1:1 hydrochloride

a) 2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-ethyl}-carbamic acid tert-butyl ester

As described for example 8, 4-(4-fluoro-benzyloxy)-phthalic acid (1.012 g, 3.5 mmol) was converted to the title compound (130 mg, 9%) [using tert-butyl N-(2-aminoethyl)-carbamate instead of N-acetylenediamine] which was obtained as a white solid. MS: $m/e = 415.4$ ($M+H^+$).

b) 2-(2-Amino-ethyl)-5-(4-fluoro-benzyloxy)-isoindole-1,3-dione 1:1 hydrochloride

A mixture of {2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-ethyl}-carbamic acid tert-butyl ester (200 mg, 0.5 mmol) and HCl in dioxane (4 N, 5 mL) was stirred at rt for 72 h. The precipitate was filtered off and washed with ether to afford the title compound (102 mg, 60%) as a white solid. MS: $m/e = 351.8$ ($M+H^+$).

Example 10

5-(4-Fluoro-benzyloxy)-2-(2-piperidin-1-yl-ethyl)-isoindole-1,3-dione

As described for example 8, 4-(4-fluoro-benzyloxy)-phthalic acid (200 mg, 0.7 mmol) was converted to the title compound (215 mg, 67%) [using 1-(2-aminoethyl)-piperidine instead of N-acetythylenediamine] which was obtained as a light yellow solid.

MS: $m/e = 383.3$ ($M+H^+$).

Example 11

5-(4-Fluoro-benzyloxy)-2-(2-morpholin-4-yl-ethyl)-isoindole-1,3-dione

As described for example 8, 4-(4-fluoro-benzyloxy)-phthalic acid (200 mg, 0.7 mmol) was converted to the title compound (179 mg, 68%) [using 1-(aminoethyl)-morpholine instead of N-acetythylenediamine] which was obtained as an off-white solid. MS: $m/e = 385.3$ ($M+H^+$).

Example 12

5-(4-Fluoro-benzyloxy)-2-(2-hydroxy-ethyl)-isoindole-1,3-dione

As described for example 8, 4-(4-fluoro-benzyloxy)-phthalic acid (200 mg, 0.7 mmol) was converted to the title compound (116 mg, 53%) [using ethanolamine instead of N-acetythylenediamine] which was obtained as a white solid.

MS: $m/e = 316.2$ ($M+H^+$).

Example 13

5-(4-Fluoro-benzyloxy)-2-(2-methoxy-ethyl)-isoindole-1,3-dione

As described for example 8, 4-(4-fluoro-benzyloxy)-phthalic acid (200 mg, 0.7 mmol) was converted to the title compound (137 mg, 60%) [using 2-methoxyethylamine instead of N-acetythylenediamine] which was obtained as a white solid.

MS: $m/e = 330.4$ ($M+H^+$).

Alternatively, 5-(4-fluoro-benzyloxy)-2-(2-methoxy-ethyl)-isoindole-1,3-dione was also prepared starting from 5-hydroxy-2-(2-methoxy-ethyl)-isoindole-1,3-dione.

a) 5-Hydroxy-2-(2-methoxy-ethyl)-isoindole-1,3-dione

A mixture of 4-hydroxyphthalic acid (0.91 mg, 5.0 mmol) and carbonyldiimidazole (0.85 g, 5 mmol) in N,N-dimethylacetamide (3 mL) was stirred at room temperature for 45 min. To this mixture was added 2-methoxyethylamine (0.47 g, 6.0 mmol) and the resulting mixture heated under reflux for 15 min. After cooling to room temperature, the mixture was evaporated and the residue was purified by chromatography (SiO₂, CH₂Cl₂: 2N NH₃/MeOH 19:1 to 9:1) to afford the title compound (1.1 g, 95%) as a light yellow solid. MS: m/e = 220.1 (M-H⁻).

b) 5-(4-Fluoro-benzyloxy)-2-(2-methoxy-ethyl)-isoindole-1,3-dione

A mixture of 5-hydroxy-2-(2-methoxy-ethyl)-isoindole-1,3-dione, from example 13a; (570 mg, 2.6 mmol), potassium carbonate (445 mg, 3.2 mmol) and 4-fluorobenzylbromide (526 mg, 2.8 mmol) in acetone (40 mL) was heated under reflux for 17 h. After cooling to room temperature, the mixture was filtered, evaporated and the residue was purified by crystallisation from diethylether : heptane (1:2) to afford the title compound (560 mg, 66%) as a white solid. MS: m/e = 330.4 (M+H⁺).

Example 14

(S)-2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-3-hydroxy-propionamide

As described for example 8, 4-(4-fluoro-benzyloxy)-phthalic acid (200 mg, 0.7 mmol) was converted to the title compound (126 mg, 51%) [using L-(-)-serinamide HCl and pyridine (65 mg, 0.8 mmol) instead of N-acetythylenediamine] which was obtained as a white solid. MS: m/e = 359.2 (M+H⁺).

Example 15

(S)-2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-propionic acid methyl ester

As described for example 14, 4-(4-fluoro-benzyloxy)-phthalic acid (200 mg, 0.7 mmol) was converted to the title compound (148 mg, 53%) [using L-alanine methyl ester-HCl instead of L-(-)-serinamide HCl] which was obtained as a white solid after crystallisation from MeOH. MS: m/e = 358.2 (M+H⁺).

Example 16

(S)-2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-3-phenyl-propionamide

As described for example 8, 4-(4-fluoro-benzyloxy)-phthalic acid (200 mg, 0.7 mmol) was converted to the title compound (221 mg, 77%) [using L-(-)-phenylalanine-
5 amide instead of N-acetythylenediamine] which was obtained as a white solid.
MS: m/e = 359.2 (M+H⁺).

Example 17

(R)-2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-propionamide

As described for example 14, 4-(4-fluoro-benzyloxy)-phthalic acid (200 mg, 0.7 mmol) was converted to the title compound (148 mg, 53%) [using H-D-Ala-NH₂ HCl
10 instead of L-(-)-serinamide HCl] which was obtained as a white solid after trituration with MeOH. MS: m/e = 343.2 (M+H⁺).

Example 18

2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-3-phenyl-propionamide

As described for example 8, 4-(4-fluoro-benzyloxy)-phthalic acid (1.2 g, 4.0 mmol) was converted to the title compound (200 mg, 13%) [using 2-methylalanine methyl ester
instead of N-acetythylenediamine] which was obtained as a white solid.
20 MS: m/e = 372.3 (M+H⁺).

Example 19

2-(2-Ethylsulfanyl-ethyl)-5-(4-fluoro-benzyloxy)-isoindole-1,3-dione

As described for example 8, 4-(4-fluoro-benzyloxy)-phthalic acid (290 mg, 1.0 mmol) was converted to the title compound (212 mg, 59%) [2-(ethylthio)ethylamine
25 instead of N-acetythylenediamine] which was obtained as a white solid after crystallisation from MeOH. MS: m/e = 360.2 (M+H⁺).

Example 20

(R,S)-2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-butyramide

As described for example 1 c, 4-(4-fluoro-benzyloxy)-phthalic acid (290 mg, 1.0 mmol) was converted to the title compound (210 mg, 59%) [using (rac)-2-aminobutyramide hydrochloride instead of glycine hydrochloride and N,N-dimethylacetamide instead of 1-methyl-2-pyrrolidinone] which was obtained as a white
30 solid. MS: m/e = 357.3 (M+H⁺).

Example 21

4-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-butyramide

As described for example 20, 4-(4-fluoro-benzyloxy)-phthalic acid (290 mg, 1.0 mmol) was converted to the title compound (103 mg, 29%) [using 4-aminobutyramide hydrochloride instead of (rac)-2-aminobutyramide hydrochloride] which was obtained as a white solid after crystallisation from MeOH. MS: m/e = 357.3 (M+H⁺).

Example 22

4-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-3-hydroxy-butyramide

As described for example 21, 4-(4-fluoro-benzyloxy)-phthalic acid (290 mg, 1.0 mmol) was converted to the title compound (108 mg, 22%) [using 4-amino-3-hydroxybutyramide hydrochloride instead of 4-aminobutyramide hydrochloride] which was obtained as an off-white solid. MS: m/e = 373.3 (M+H⁺).

Example 23

5-(3-Fluoro-benzyloxy)-2-(2-methoxy-ethyl)-isoindole-1,3-dione

As described for example 13b, a mixture of 5-hydroxy-2-(2-methoxy-ethyl)-isoindole-1,3-dione (example 13a, 400 mg, 1.8 mmol) was converted to title compound (427 mg, 72%) (using 3-fluorobenzylbromide instead of 4-fluorobenzylbromide) (376 mg, 2.0 mmol) and heating under reflux for 67 h which was obtained as a white solid after purification by chromatography (SiO₂, CH₂Cl₂: 2N NH₃/MeOH 99:1 to 19:1). MS: m/e = 330.4 (M+H⁺).

Example 24

(S)-5-(4-Fluoro-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione

As described for example 8, 4-(4-fluoro-benzyloxy)-phthalic acid (300 mg, 1.0 mmol) was converted to the title compound (286 mg, 81%) [using (S)-1-methoxy-2-propylamine instead of N-acetylenehtylenediamine] which was obtained as a white solid after trituration with hexane. MS: m/e = 344.4 (M+H⁺).

Example 25

(S)-5-(3-Fluoro-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dionea) (S)-5-Hydroxy-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione

As described for example 13a, 4-hydroxyphthalic acid (8.0 g, 44 mmol) was converted to the title compound (8.7 g, 84%) [using (S)-1-methoxy-2-propylamine instead of 2-methoxyethylamine] which was obtained as light-yellow crystals after purification by chromatography (SiO₂, EtOAc:Hexane 1:1). MS: m/e = 234.1 (M-H⁻).

b) (S)-5-(3-Fluoro-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione

As described for example 23, a mixture of (S)-5-hydroxy-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione (example 25a, 200 mg, 0.9 mmol) was converted to title compound (200 mg, 69%) (using 3-fluorobenzylbromide instead of 4-fluorobenzylbromide), after heating under reflux for 1 h, which was obtained as a white solid after crystallisation from ethylacetate : hexane (1 : 1). MS: $m/e = 343.2$ (M^+).

Example 26

(S)-5-(2-Fluoro-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione

As described for example 23, a mixture of (S)-5-hydroxy-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione (example 25a, 200 mg, 0.9 mmol) was converted to the title compound (235 mg, 81%) (using 2-fluorobenzylbromide instead of 4-fluorobenzylbromide), after heating under reflux for 1 h, which was obtained as a white solid after crystallisation from ethylacetate : hexane (1 : 1). MS: $m/e = 343.2$ (M^+).

Example 27

15 (S)-2-(2-Methoxy-1-methyl-ethyl)-5-(4-trifluoromethyl-benzyloxy)-isoindole-1,3-dione

As described for example 23, a mixture of (S)-5-hydroxy-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione (example 25a, 200 mg, 0.9 mmol) was converted to the title compound (265 mg, 79%) [using 4-(trifluoromethyl)benzylbromide instead of 4-fluorobenzylbromide], after heating under reflux for 1.5 h, which was obtained as a white solid after crystallisation from ethylacetate : hexane (1 : 1). MS: $m/e = 393.2$ (M^+).

Example 28

(S)-5-(4-Bromo-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione

As described for example 23, a mixture of (S)-5-hydroxy-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione (example 25a, 200 mg, 0.9 mmol) was converted to the title compound (296 mg, 86%) (using 4-bromobenzylbromide instead of 4-fluorobenzylbromide), after heating under reflux for 1.5 h, which was obtained as a white solid after crystallisation from ethylacetate : hexane (1 : 1). MS: $m/e = 403.2/405.2$ (M^+).

Example 29

(S)-5-(3-Methoxy-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione

30 As described for example 23, a mixture of (S)-5-hydroxy-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione (example 25a, 200 mg, 0.9 mmol) was converted to the title compound (280 mg, 93%) (using 3-methoxybenzylbromide instead of 4-fluorobenzylbromide), after heating under reflux for 3 h, which was obtained as a white solid after crystallisation from ethylacetate : hexane (1 : 1). MS: $m/e = 355.4$ (M^+).

Example 30

(S)-4-[2-(2-Methoxy-1-methyl-ethyl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-ylloxymethyl]-benzonitrile

As described for example 23, a mixture of (S)-5-hydroxy-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione (example 25a, 200 mg, 0.9 mmol) was converted to the title compound (280 mg, 94%) (using 3-bromomethyl benzonitrile instead of 4-fluorobenzylbromide), after heating under reflux for 1.5 h, which was obtained as a white solid after crystallisation from ethylacetate : hexane (1 : 1). MS: m/e = 350.2 (M⁺).

Example 31

10 (S)-5-(3,4-Difluoro-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione

As described for example 23, a mixture of (S)-5-hydroxy-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione (example 25a, 200 mg, 0.9 mmol) was converted to title compound (235 mg, 77%) (using alpha-bromo-3,4-difluorotoluene instead of 4-fluorobenzylbromide), after heating under reflux for 2 h, which was obtained as a white solid after crystallisation from ethylacetate : hexane (1 : 1). MS: m/e = 361.2 (MH).

Example 32

(S)-2-(2-Methoxy-1-methyl-ethyl)-5-(4-trifluoromethoxy-benzyloxy)-isoindole-1,3-dione

As described for example 23, a mixture of (S)-5-hydroxy-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione (example 25a, 200 mg, 0.9 mmol) was converted to the title compound (306 mg, 88%) [using (4-trifluoromethoxy)benzyl bromide instead of 4-fluorobenzylbromide], after heating under reflux for 3 h, which was obtained as a white solid after crystallisation from ethylacetate : hexane (1 : 1). MS: m/e = 409.4 (M+H⁺).

Example 33

25 5-(4-Fluoro-benzyloxy)-2-(tetrahydro-furan-2-ylmethyl)-isoindole-1,3-dione

As described for example 1d, 4-(4-fluoro-benzyloxy)-phthalic acid (290 mg, 1.0 mmol) was converted to the title compound (249 mg, 70%) [using tetrahydrofurfurylamine instead of glycineamide hydrochloride and pyridine; and N,N-dimethylacetamide instead of 1-methyl-2-pyrrolidinone] which was obtained as a white solid.

30 MS: m/e = 356.3 (M+H⁺).

Example 34

5-(4-Fluoro-benzyloxy)-2-thiophen-2-ylmethyl-isoindole-1,3-dione

As described for example 33, 4-(4-fluoro-benzyloxy)-phthalic acid (290 mg, 1.0 mmol) was converted to the title compound (269 mg, 73%) [using 2-thiophenemethyl-

amine instead of tetrahydrofurfurylamine] which was obtained as a white solid.

MS: $m/e = 367.1$ ($M+H^+$).

Examples 35

(S)-2-[5-(4-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-thiopropionamide

5 A mixture of (S)-2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-propionamide (171 mg, 0.5 mmol) and Lawesson's reagent (243 mg, 0.6 mmol) in tetrahydrofuran (20 mL) was stirred at room temperature for 72 h. Then the mixture was evaporated and the residue was purified by chromatography (SiO_2 , CH_2Cl_2 : 2N NH_3 /MeOH 99:1 to 19:1) to afford the title compound (65 mg, 36%) as an off-white
10 solid after recrystallisation from MeOH. MS: $m/e = 359.2$ ($M+H^+$).

Example 36

5-(4-Fluoro-benzyloxy)-2-piperidin-4-yl-isoindole-1,3-dione 1:1 hydrochloride

As described for example 33, 4-(4-fluoro-benzyloxy)-phthalic acid (590 mg, 2.0 mmol) was converted to the title compound [using using 4-amino-1-Boc-piperidine
15 instead of tetrahydrofurfurylamine] followed by treatment with HCl in dioxane (4 N, 2 mL) and stirring at room temperature for 12 h. The precipitate was filtered off and washed with acetone and ether to afford the title compound (73 mg, 9%) as a white solid. MS: $m/e = 355.3$ ($M+H^+$).

Example 37

5-(4-Fluoro-benzyloxy)-2-(5-methyl-isoxazol-3-yl)-isoindole-1,3-dione

As described for example 33, 4-(4-fluoro-benzyloxy)-phthalic acid (290 mg, 1.0 mmol) was converted to the title compound (166 mg, 47%) [using 3-amino-5-methylisoxazole instead of tetrahydrofurfurylamine] as a white solid.
MS: $m/e = 353.3$ ($M+H^+$).

25

Example 38

2-(2-Ethanesulfinyl-ethyl)-5-(4-fluoro-benzyloxy)-isoindole-1,3-dione

A mixture of 2-(2-ethylsulfinyl-ethyl)-5-(4-fluoro-benzyloxy)-isoindole-1,3-dione (96 mg, 0.27 mmol) and 3-phenyl-2-(phenylsulfonyl)oxaziridine (84 mg, 0.32 mmol) in CH_2Cl_2 (3 mL) was stirred at room temperature for 72 h. Then the mixture was
30 evaporated and the residue was purified by chromatography (SiO_2 , CH_2Cl_2 : 2N NH_3 /MeOH 97:3 to 9:1) to afford the title compound (34 mg, 34%) as a white solid. MS: $m/e = 375.2$ ($M+H^+$).

Example 39

(S)-2-[2-(4-Fluoro-benzyloxy)-5,7-dioxo-5,7-dihydro-pyrrolo[3,4-b]pyridin-6-yl]-
propionamide

a) 1-Oxy-pyridine-2,3-dicarboxylic acid diethyl ester

- 5 A mixture of diethyl 2,3-pyridinedicarboxylate (10 g, 45 mmol) and 3-chloroperoxy-benzoic acid (13.8 g, 56 mmol) in CH_2Cl_2 (150 mL) was stirred at 0 °C and allowed to warm up room temperature over 30 min and left overnight. Then half-evaporation gave a residue which was purified by chromatography (SiO_2 , EtOAc: EtOAc:MeOH, 19:1) to afford the title compound (8.2 g, 77%) as a white solid. MS: $m/e = 239.2$ (M^+)

10 b) 6-Chloro-pyridine-2,3-dicarboxylic acid diethyl ester

A mixture of 1-oxy-pyridine-2,3-dicarboxylic acid diethyl ester (8.0 g, 33 mmol) and phosphoryloxychloride (25.6 g, 167 mmol) was heated at 100 °C for 1.5 h. After cooling to room temperature, the mixture was evaporated and then diluted with CH_2Cl_2 . Then the organic phase was washed with water, brine and then dried over sodium sulfate.

- 15 Filtration and evaporation gave a residue which was purified by chromatography (SiO_2 , heptane-EtOAc 9:1 to 1:1) to afford the title compound (5.5 g, 63%) as a colourless oil. MS: $m/e = 258.1$ ($\text{M}+\text{H}^+$).

c) 6-(4-Fluoro-benzyloxy)-pyridine-2,3-dicarboxylic acid diethyl ester

- A mixture of sodium hydride (55% in mineral oil, 254 mg, 0.6 mmol) and 4-fluorobenzylalcohol in tetrahydrofuran (30 mL) was heated at 60 °C for 30 min and then after cooling at 0 °C a solution of 6-chloro-pyridine-2,3-dicarboxylic acid diethyl ester (1.3 g, 0.5 mmol) in tetrahydrofuran (20 mL) was added over 10 min. The resulting mixture was kept at 0 °C for 1 h and then water was added. The mixture was then extracted with ethylacetate and the combined organic extracts washed with water, brine and then dried over sodium sulfate. Filtration and evaporation gave a residue which was purified by chromatography (SiO_2 , heptane-EtOAc 9:1 to 3:1) to afford the title compound (1.3 g, 74%) as a colourless oil. MS: $m/e = 348.4$ ($\text{M}+\text{H}^+$).

d) 6-(4-Fluoro-benzyloxy)-pyridine-2,3-dicarboxylic acid diethyl ester

- A mixture of 6-(4-fluoro-benzyloxy)-pyridine-2,3-dicarboxylic acid diethyl ester (1.2 g, 3.5 mmol) and potassium hydroxide (0.6 g, 11.5 mmol) in water (25 mL) and tetrahydrofuran (10 mL) was heated at 80 °C for 4 h. After cooling to room temperature, the mixture was washed with ether. The aqueous phase was then acidified to pH 2 at 0 °C with concentrated HCl and then extracted with ethyl acetate. The organic extracts were then washed with water and brine and dried over sodium sulfate. Filtration and evaporation gave a residue which was purified by crystallisation from diethylether :

heptane (2:1) to afford the title compound (0.9 g, 88%) as a white solid.

MS: $m/e = 290.0$ ($M-H^+$).

e) (S)-2-[2-(4-Fluoro-benzyloxy)-5,7-dioxo-5,7-dihydro-pyrrolo[3,4-b]pyridin-6-yl]-propionamide

- 5 As described for example 14, 4-(4-fluoro-benzyloxy)-phthalic acid (200 mg, 0.7 mmol) was converted to the title compound (40 mg, 12%) [using H-Ala-NH₂ HCl instead of L-(-)-serinamide HCl] which was obtained as a white solid after purification by HPLC (Waters Xterra RP18 (5 μ m x 50 x 19mm) eluting with AcCN/ 0.1%TFA/Water. MS: $m/e = 344.3$ ($M+H^+$).

10 **Example 40**

5-(4-Fluoro-benzyloxy)-2-(3-morpholin-4-yl-propyl)-isoindole-1,3-dione

a) 5-Hydroxy-2-(3-morpholin-4-yl-propyl)-isoindole-1,3-dione

- 15 As described for example 13a, 4-hydroxyphthalic acid (1.0 g, 5.5 mmol) was converted to the title compound (780 mg, 49%) [using 4-(3-aminopropyl)morpholine instead of 2-methoxyethylamine] which was obtained as a light yellow solid. MS: $m/e = 291.3$ ($M+H^+$).

b) 5-(4-Fluoro-benzyloxy)-2-(3-morpholin-4-yl-propyl)-isoindole-1,3-dione

- 20 As described for example 13b, 5-hydroxy-2-(3-morpholin-4-yl-propyl)-isoindole-1,3-dione (770 mg, 2.65 mmol) was converted to the title compound (210 mg, 28%) which was obtained as a white solid after purification by chromatography (SiO₂, CH₂Cl₂: 2N NH₃/MeOH 99:1 to 19:1). MS: $m/e = 399.4$ ($M+H^+$).

Example 41

2-(4-Fluoro-benzyl)-5-(4-fluoro-benzyloxy)-isoindole-1,3-dione

- 25 A mixture of 5-hydroxy-isoindole-1,3-dione (200 mg, 1.0 mmol), potassium carbonate (178 mg, 1.05 mmol), 4-fluorobenzylbromide (204 mg, 1.05 mmol) in ethanol (5 mL) was heated at 80 °C overnight. After cooling to room temperature, the reaction mixture was filtered and evaporated. The residue was purified by chromatography (SiO₂, heptane-CH₂Cl₂ 2:3 then CH₂Cl₂- 2N NH₃-MeOH 99:1 to 95:5) to afford the title compound (13 mg, 3%) as a white solid. MS: $m/e = 379.4$ (M^+).

30 **Example 42**

5-(4-Fluoro-benzyloxy)-2-(3,3,3-trifluoro-2-hydroxy-propyl)-isoindole-1,3-dione

- As described for example 8, 4-(4-fluoro-benzyloxy)-phthalic acid (500 mg, 1.7 mmol) was converted to the title compound (567 mg, 86%) [using 3-amino-1,1,1-trifluoro-2-propanol instead of N-acetythylenediamine] which was obtained as a white

solid after purification by chromatography (SiO₂, hexane-EtOAc 2:1). MS: m/e = 384.3 (M⁺).

Example 43

(5-(4-Fluoro-benzyloxy)-2-(3,3,3-trifluoro-2-methoxy-propyl)-isoindole-1,3-dione

- 5 To a suspension of sodium hydride (45 mg, 1.0 mmol) in tetrahydrofuran (8 mL) was added 5-(4-fluoro-benzyloxy)-2-(3,3,3-trifluoro-2-hydroxy-propyl)-isoindole-1,3-dione (360 mg, 0.9 mmol) at room temperature and then iodomethane (159 mg, 1.1 mmol) was added and then after 14 h, water was added. The mixture was then extracted with ethylacetate and the combined organic extracts washed with water, brine and then
10 dried over sodium sulfate. Filtration and evaporation gave a residue which was purified by chromatography (SiO₂, hexane-EtOAc 3:1) to afford the title compound (120 mg, 32%) as a white solid. MS: m/e = 398.4 (M+H⁺).

Example 44

(S)-2-[5-(3-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-propionamide

- 15 a) (S)-2-(5-Hydroxy-1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propionamide

As described for example 13a, 4-hydroxyphthalic acid (1.82 g, 1.0 mmol) was converted to the title compound (845 mg, 36%) [using H-Ala-NH₂ HCl and pyridine instead of 2-methoxyethylamine] which was obtained as a light yellow solid. MS: m/e = 233.0 (M-H⁻).

- 20 b) (S)-2-[5-(3-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-propionamide

As described for example 13b, (S)-2-(5-hydroxy-1,3-dioxo-1,3-dihydro-isoindol-2-yl)-propionamide (250 mg, 1.1 mmol) was converted to the title compound (70 mg, 19%) (using 3-fluorobenzylbromide instead of 4-fluorobenzylbromide) which was obtained as a white solid after purification by chromatography (SiO₂, CH₂Cl₂: 2N NH₃/MeOH
25 9:1). MS: m/e = 343.3 (M+H⁺).

Example 45

(2-[5-(3-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetamide

- a) 4-(3-Fluoro-benzyloxy)-phthalic acid

30 As described for example 1a and b, 4-hydroxyphthalic acid (5.0 g, 27 mmol) was converted to the title compound (6.9 g, 87%) [using 3-fluorobenzylbromide instead of 4-fluorobenzylbromide] which was obtained as a white solid after recrystallisation from diethyl ether : hexane (1:3). MS: m/e = 289.0 (M-H⁻).

b) (2-[5-(3-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetamide

As described for example 1d, 4-(3-fluoro-benzyloxy)-phthalic acid (200 mg, 0.69 mmol) was converted to the title compound (151 mg, 66%) which was obtained as a white solid. MS: $m/e = 328.2$ (M^+).

5

Example 46

5-(3-Fluoro-benzyloxy)-2-(2-hydroxy-ethyl)-isoindole-1,3-dione

As described for example 8, 4-(3-fluoro-benzyloxy)-phthalic acid (200 mg, 0.69 mmol) was converted to the title compound (179 mg, 82%) [using ethanolamine instead of N-acetylenediamine] which was obtained as a white solid after purification by chromatography (SiO_2 , EtOAc: Hexane 1:1). MS: $m/e = 315.2$ (M^+).

10

Example 47

2-(2-Ethylsulfanyl-ethyl)-5-(3-fluoro-benzyloxy)-isoindole-1,3-dione

As described for example 8, 4-(3-fluoro-benzyloxy)-phthalic acid (200 mg, 0.69 mmol) was converted to the title compound (217 mg, 88%) [2-(ethylthio)ethylamine instead of N-acetylenediamine] which was obtained as an off-white solid after purification by chromatography (SiO_2 , EtOAc: Hexane 1:3). MS: $m/e = 359.2$ (M^+).

15

Example 48

(5-(3-Fluoro-benzyloxy)-2-(3,3,3-trifluoro-2-hydroxy-propyl)-isoindole-1,3-dione

As described for example 8, 4-(3-fluoro-benzyloxy)-phthalic acid (200 mg, 0.69 mmol) was converted to the title compound (235 mg, 89%) [using 3-amino-1,1,1-trifluoro-2-propanol instead of N-acetylenediamine] which was obtained as a white solid after purification by chromatography (SiO_2 , hexane-EtOAc 2:1). MS: $m/e = 383.2$ (M^+).

20

Example 49

25 5-(3-Fluoro-benzyloxy)-2-(3,3,3-trifluoro-2-methoxy-propyl)-isoindole-1,3-dione

As described for example 43, (5-(3-fluoro-benzyloxy)-2-(3,3,3-trifluoro-2-hydroxy-propyl)-isoindole-1,3-dione (120 mg, 0.31 mmol) was converted to the title compound (25 mg, 21%) which was obtained as a white solid after purification by chromatography (SiO_2 , hexane-EtOAc 3:1). MS: $m/e = 398.3$ ($M+H^+$).

Example 50

2-(2-Amino-ethyl)-5-(3-fluoro-benzyloxy)-isoindole-1,3-dione 1:1 hydrochloridea) {2-[5-(3-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-ethyl}-carbamic acid tert-butyl ester

5 As described for example 9a, 4-(3-fluoro-benzyloxy)-phthalic acid (300 mg, 1.0 mmol) was converted to the title compound (338 mg, 79%) which was obtained as a white solid after purification by chromatography (SiO₂, hexane-EtOAc 3:1). MS: m/e = 415.4 (M+H⁺).

b) 2-(2-Amino-ethyl)-5-(3-fluoro-benzyloxy)-isoindole-1,3-dione 1:1 hydrochloride

10 As described for example 9b, {2-[5-(3-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-ethyl}-carbamic acid tert-butyl ester (311 mg, 0.75 mmol) was converted to the title compound (249 mg, 94%) which was obtained as a white solid. MS: m/e = 315.3 (M+H⁺).

Example 51

15 [5-(3-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetonitrile

As described for example 1d, 4-(3-fluoro-benzyloxy)-phthalic acid (200 mg, 0.69 mmol) was converted to the title compound (176 mg, 82%) (using aminoacetonitrile instead of glycine hydrochloride) which was obtained as a white solid after purification by chromatography (SiO₂, hexane-EtOAc 3:1). MS: m/e = 310.2 (M⁺).

20

Example 52

(S)-2-[5-(3-Fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-3-hydroxy-propionamide

As described for example 8, 4-(3-fluoro-benzyloxy)-phthalic acid (200 mg, 0.69 mmol) was converted to the title compound (76 mg, 31%) [using L-(-)-serinamide HCl and pyridine (65 mg, 0.8 mmol) instead of N-acetylenediamine] which was obtained
25 as a light-orange solid. MS: m/e = 359.2 (M+H⁺).

Example 53

(S)-5-(3,5-Bis-trifluoromethyl-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione

30 As described for example 23, a mixture of (S)-5-hydroxy-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione (200 mg, 0.9 mmol) was converted to title compound (314 mg, 81%) [using 3,5-bis-(trifluoromethyl)benzyl bromide instead of 4-fluorobenzyl-bromide], after heating under reflux for 1 h, which was obtained as a white solid after crystallisation from ethylacetate : hexane (1 : 1). MS: m/e = 461.2 (M⁺).

Example A

Tablets of the following composition are produced in a conventional manner:

	<u>mg/Tablet</u>
Active ingredient	100
5 Powdered lactose	95
White corn starch	35
Polyvinylpyrrolidone	8
Na carboxymethylstarch	10
Magnesium stearate	2
10 Tablet weight	<u>250</u>

Example B

Tablets of the following composition are produced in a conventional manner:

	<u>mg/Tablet</u>
Active ingredient	200
15 Powdered lactose	100
White corn starch	64
Polyvinylpyrrolidone	12
Na carboxymethylstarch	20
Magnesium stearate	4
20 Tablet weight	<u>400</u>

Example C

Capsules of the following composition are produced:

	<u>mg/Capsule</u>
Active ingredient	50
5 Crystalline lactose	60
Microcrystalline cellulose	34
Talc	5
Magnesium stearate	1
Capsule fill weight	<u>150</u>

10 The active ingredient having a suitable particle size, the crystalline lactose and the microcrystalline cellulose are homogeneously mixed with one another, sieved and thereafter talc and magnesium stearate are admixed. The final mixture is filled into hard gelatine capsules of suitable size.


Example D

- 15 An injection solution may have the following composition and is manufactured in usual manner:

Active substance	1.0 mg
1 N HCl	20.0 µl
acetic acid	0.5 mg
20 NaCl	8.0 mg
phenol	10.0 mg
1 N NaOH	q.s. ad pH 5
H ₂ O	q.s. ad 1 ml

Claims

1. Compounds of the general formula



I

wherein

- 5 X is -N= or -CH=;
- R¹ is -CO-NR⁵R⁶; -CHR⁷-(CH₂)_n-CO-NR⁵R⁶;
 -(CH₂)_n-NR⁵R⁶;
 -(CH₂)_n-COOR⁸;
 -(CH₂)_n-CN;
 10 -CHR⁷-(CH₂)_n-CF₃;
 -(CH₂)_n-NH-COR⁹;
 -(CH₂)_n-NH-COOR⁸;
 -(CH₂)_n-piperidinyl, -(CH₂)_n-morpholinyl, -(CH₂)_n-tetrahydrofuranyl;
 -(CH₂)_n-thiophenyl or -(CH₂)_n-isoxazolyl, wherein the heterocyclic ring may
 15 be substituted by C₁-C₆-alkyl;
 -(CH₂)_n-phenyl, wherein the phenyl ring may be substituted by halogen or
 halogen-(C₁-C₆)-alkyl;
 -(CH₂)_p-OR⁸;
 -(CH₂)_p-SR⁸;
 20 -(CH₂)_p-SO-R⁹; or
 -(CH₂)_n-CS-NR⁵R⁶;
- R² is hydrogen; C₁-C₆-alkyl;
 -(CH₂)_p-OR¹⁰;
 -(CH₂)_p-SR¹⁰; or benzyl;
- 25 R³ is hydrogen or C₁-C₆-alkyl;
- R⁴ is halogen, halogen-(C₁-C₆)-alkyl, cyano, C₁-C₆-alkoxy or
 halogen-(C₁-C₆)-alkoxy;
- R⁵ and R⁶ are independently from each other hydrogen or C₁-C₆-alkyl;
- R⁷ is hydrogen, hydroxy or C₁-C₆-alkoxy;

R^8 is hydrogen or C_1 - C_6 -alkyl;

R^9 is C_1 - C_6 -alkyl;

R^{10} is hydrogen or C_1 - C_6 -alkyl;

m is 1, 2 or 3;

5 n is 0, 1 or 2; and

p is 1 or 2;

as well as their pharmaceutically acceptable salts.

2. Compounds of formula I according to claim 1, wherein X is $-\text{CH}=\text{}$.

3. Compounds of formula I according to claim 1 or 2, wherein R^1 is $-\text{CO}-\text{NR}^5\text{R}^6$ or
10 $-\text{CHR}^7-(\text{CH}_2)_n-\text{CO}-\text{NR}^5\text{R}^6$, and wherein R^5 and R^6 are independently from each other hydrogen or C_1 - C_6 -alkyl, R^7 is hydrogen, hydroxy or C_1 - C_6 -alkoxy and n is 0, 1 or 2.

4. Compounds of formula I according to claim 3, which compounds are selected from the group consisting of

2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetamide,
15 (S)-2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-propionamide,
(S)-2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-3-hydroxy-propionamide,
(R)-2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-propionamide,
2-[5-(3-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-propionamide,
20 (2-[5-(3-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetamide, and
2-[5-(3-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-3-hydroxy-propionamide.

5. Compounds of formula I according to claim 1 or 2, wherein R^1 is $-(\text{CH}_2)_n-\text{NR}^5\text{R}^6$, $-(\text{CH}_2)_n-\text{NH}-\text{COR}^9$ or $-(\text{CH}_2)_n$ -piperidiny, and wherein R^5 and R^6 are
25 independently from each other hydrogen or C_1 - C_6 -alkyl, R^9 is C_1 - C_6 -alkyl and n is 0, 1, or 2.

6. Compounds of formula I according to claim 5, which compounds are selected from the group consisting of

N-{2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-ethyl}-acetamide,
30 2-(2-amino-ethyl)-5-(4-fluoro-benzyloxy)-isoindole-1,3-dione, and
5-(4-fluoro-benzyloxy)-2-piperidin-4-yl-isoindole-1,3-dione.

7. Compounds of formula I according to claim 1 or 2, wherein R^1 is $-(CH_2)_p-OR^8$ or $-CHR^7-(CH_2)_n-CF_3$, and wherein R^8 is hydrogen, hydroxy or C_1-C_6 -alkoxy, R^7 is hydrogen or C_1-C_6 -alkyl and p is 1 or 2.

8. Compounds of formula I according to claim 7, which compounds are selected from the group consisting of

5 5-(4-fluoro-benzyloxy)-2-(2-hydroxy-ethyl)-isoindole-1,3-dione,
5-(4-fluoro-benzyloxy)-2-(2-methoxy-ethyl)-isoindole-1,3-dione,
5-(3-fluoro-benzyloxy)-2-(2-methoxy-ethyl)-isoindole-1,3-dione,
(S)-5-(4-fluoro-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione,
10 (S)-5-(3-fluoro-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione,
(S)-5-(2-fluoro-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione,
(S)-2-(2-methoxy-1-methyl-ethyl)-5-(4-trifluoromethyl-benzyloxy)-isoindole-1,3-dione,
(S)-5-(4-bromo-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione,
(S)-5-(3,4-difluoro-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione,
15 5-(3-fluoro-benzyloxy)-2-(2-hydroxy-ethyl)-isoindole-1,3-dione,
5-(4-fluoro-benzyloxy)-2-(3,3,3-trifluoro-2-hydroxy-propyl)-isoindole-1,3-dione, and
5-(3,5-bis-trifluoromethyl-benzyloxy)-2-(2-methoxy-1-methyl-ethyl)-isoindole-1,3-dione.

9. Compounds of formula I according to claim 1 or 2, wherein R^1 is $-(CH_2)_p-SR^8$; $-(CH_2)_p-SO-R^9$; or $-(CH_2)_n-CS-NR^5R^6$, and wherein R^5 and R^6 are independently from each other hydrogen or C_1-C_6 -alkyl, R^8 is hydrogen or C_1-C_6 -alkyl, R^9 is C_1-C_6 -alkyl and n is 0, 1, or 2.

10. Compounds of formula I according to claim 9, which compounds are selected from the group consisting of

25 2-(2-ethylsulfanyl-ethyl)-5-(4-fluoro-benzyloxy)-isoindole-1,3-dione,
(S)-2-[5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-thiopropionamide,
and
2-(2-ethylsulfanyl-ethyl)-5-(3-fluoro-benzyloxy)-isoindole-1,3-dione.

11. Compounds of formula I according to claim 1 or 2, wherein R^1 is $-(CH_2)_n-CN$ and n is 0, 1 or 2.


12. Compounds of formula I according to claim 11, which compounds are selected from the group consisting of

5-(4-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetonitrile, and
[5-(3-fluoro-benzyloxy)-1,3-dioxo-1,3-dihydro-isoindol-2-yl]-acetonitrile.

13. Compounds of formula I according to claim 1, wherein R^4 is halogen.


14. A process for the manufacture of a compound of formula I according to any one of claims 1 to 13 as well as its pharmaceutically acceptable salt, which process comprises

a) reacting a compound of formula




II

with a compound of formula



III


to obtain a compound of formula



I

and, if desired, converting a compound of formula I into a pharmaceutically acceptable salt; or


b) reacting a compound of formula



IV


- 37 -

with a compound of formula



V

to obtain a compound of formula



I

- 5 and, if desired, converting a compound of formula I into a pharmaceutically acceptable salt.
15. A compound of formula I according to any one of claims 1 to 13, when manufactured by a process according to claim 14.
16. A medicament containing one or more compounds as claimed in any one of
 - 10 claims 1 to 13 and pharmaceutically acceptable excipients for the treatment and prevention of diseases which are mediated by monoamine oxidase B inhibitors.
17. A medicament containing one or more compounds as claimed in any one of claims 1 to 13 and pharmaceutically acceptable excipients for the treatment and prevention of Alzheimer's disease and senile dementia.
- 15 18. A compound of formula I according to any one of claims 1 to 13 as well as its pharmaceutically acceptable salts for the treatment or prevention of diseases.
19. The use of a compound of formula I according to any one of claims 1 to 13 as well as its pharmaceutically acceptable salts for the manufacture of medicaments for the treatment and prevention of diseases which are mediated by monoamine oxidase B
 - 20 inhibitors.
20. The use according to claim 19, wherein the disease is Alzheimer's disease or senile dementia.
21. The invention as herein before described.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/02931

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D209/48 C07D405/06 C07D409/06 C07D413/06 C07D471/04
A61K31/4035 A61K31/437 A61P25/28 //(C07D471/04,221:00,
209:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, EMBASE, BIOSIS, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 504 574 A (WAKAMOTO PHARMA CO LTD) 23 September 1992 (1992-09-23) abstract; example 10	1-20
A	US 4 524 206 A (NEW JAMES S ET AL) 18 June 1985 (1985-06-18) abstract; examples 8,31-38	1-20

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

16 July 2003

Date of mailing of the international search report

23/07/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

vanVoorst tot Voorst, M

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 21

Claim 21 not searched in accordance with Article 6 PCT and Rule 6.2(a) PCT.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 03/02931

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 21
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet (1)) (July 1998)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/02931

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0504574	A	23-09-1992	JP 4295470 A	20-10-1992
			CA 2060569 A1	23-09-1992
			DE 69208490 D1	04-04-1996
			DE 69208490 T2	11-07-1996
			EP 0504574 A1	23-09-1992
			US 5356916 A	18-10-1994
US 4524206	A	18-06-1985	AT 387773 B	10-03-1989
			AT 291584 A	15-08-1988
			AU 581858 B2	09-03-1989
			AU 3287084 A	30-05-1985
			BE 900555 A1	11-03-1985
			CA 1285564 C	02-07-1991
			CH 660484 A5	30-04-1987
			CS 8406721 A2	17-12-1987
			CS 8507792 A2	17-12-1987
			CY 1538 A	16-11-1990
			DD 224593 A5	10-07-1985
			DE 3433327 A1	28-03-1985
			DK 171990 B1	08-09-1997
			ES 8605797 A1	16-09-1986
			FI 843522 A ,B,	13-03-1985
			FR 2555585 A1	31-05-1985
			GB 2146333 A ,B	17-04-1985
			GR 80320 A1	14-01-1985
			HK 84590 A	25-10-1990
			HU 36116 A2	28-08-1985
			IE 58020 B1	30-06-1993
			IL 72854 A	31-03-1988
			IT 1196250 B	16-11-1988
			JP 1922135 C	07-04-1995
			JP 6047586 B	22-06-1994
			JP 60084282 A	13-05-1985
			KR 8900566 B1	21-03-1989
			LU 85537 A1	29-04-1985
			NL 8402769 A	01-04-1985
			NO 843579 A ,B	13-03-1985
			NZ 209480 A	08-01-1988
			OA 7809 A	20-11-1986
			PT 79187 A ,B	01-10-1984
			SE 463368 B	12-11-1990
			SE 8404552 A	13-03-1985
			SU 1384199 A3	23-03-1988
			YU 156384 A1	28-02-1987
			YU 210386 A1	31-12-1987
			ZA 8407065 A	24-04-1985